

Assignment: HW6 [40 points]

Assigned: 2016/11/18

Due: 2016/11/29

P6.1 [5 points]

For a one-dimensional system with the Hamiltonian

$$H = \frac{p^2}{2} - \frac{1}{2q^2}, \quad (1)$$

show that there is a constant of the motion

$$D = \frac{pq}{2} - Ht. \quad (2)$$

P6.2 [5 + 3 + 3 = 11 points]

Consider a particle of mass m moving in two dimensions in a potential well. Let us choose the origin of our coordinate system at the minimum of this well. The well would be termed *isotropic* if the potential did not depend on the polar angle.

- (a) First, consider the anisotropic potential in a given Cartesian coordinate system:

$$V(x_1, x_2) = \frac{k}{2}(x_1^2 + x_2^2) + k'x_1x_2; \quad k > k' > 0. \quad (3)$$

Find the eigenfrequencies and normal modes, preferably by reasoning rather than brute-force matrix diagonalization. Give a physical interpretation of the normal modes.

- (b) Use a qualitative physics-based argument to write down two independent constants of the motion. Verify your choice using the Poisson bracket equation

$$\dot{u} = \{u, H\}_{\text{PB}} + \frac{\partial u}{\partial t}, \quad (4)$$

where $u = u(q, p, t)$ and H is the Hamiltonian.

- (c) The oscillator becomes isotropic if $k' = 0$. Again use a qualitative physics-based argument to write down an additional independent constant of motion if $k' = 0$, and verify your choice with the PB equation above.

P6.3 [8 points]

Verify the Poisson bracket equation

$$\{L_i, L_j\} = \epsilon_{ijk}L_k \quad (5)$$

among the Cartesian components of angular momentum of a spherical pendulum (i.e. with two angular degrees of freedom) of mass m in a gravitational field of acceleration \vec{g} pointing opposite to the pole. ϵ_{ijk} represents

the Levi-Civita tensor¹.

Hint: Start with expressing the Lagrangian in spherical coordinates: $\mathcal{L} = \mathcal{L}(\theta, \phi, \dot{\theta}, \dot{\phi})$.

P6.4 [2 + 5 + 2 + 2 = 11 points]

Consider a system with a time-dependent Hamiltonian

$$H(q, p, t) = H_0(q, p) - \epsilon q \sin(\omega t), \quad (6)$$

where ϵ and ω are known constants and $\frac{\partial H_0}{\partial t} = 0$.

- Derive Hamilton's canonical equations of motion for the system.
- Use a canonical transformation generating function $G(q, P, t)$ to find a new Hamiltonian H' and new canonical variables Q, P such that $H'(Q, P) = H_0(q, p)$.
Hint: The partial differential equations do not tell us how q and P are related in the generating function. We can take an educated guess though. $G = qP - \frac{\epsilon q}{\omega} \cos(\omega t)$ works.
- Verify that Hamilton's canonical equations of motion are invariant under the transformation.
- Suggest a possible physical interpretation of the time-dependent term in H .

P6.5 [2 + 3 = 5 points]

The Hamiltonian $H = \frac{p^2}{2m} + \frac{m\omega^2 q^2}{2}$ describes a simple harmonic oscillator of mass m and frequency ω . Introducing the transformation

$$x_1 \equiv \omega\sqrt{m}q, \quad x_2 \equiv \frac{p}{\sqrt{m}}, \quad \tau \equiv \omega t, \quad (7)$$

we obtain $H = \frac{1}{2}(x_1^2 + x_2^2)$.

- What is the generating function $\hat{\Phi}_1(x_1, y_1)$ for the canonical transformation $\{x_1, x_2\} \rightarrow \{y_1, y_2\}$ that corresponds to the function $\Phi(q, Q) = \frac{m\omega q^2}{2} \cot Q$?
- Calculate the matrix $M_{ij} \equiv \frac{\partial x_i}{\partial y_j}$ and confirm that $\det \mathbf{M} = 1$ and $\mathbf{M}^T \epsilon \mathbf{M} = \epsilon$ (ϵ is the antisymmetric matrix used in the lectures to put the coordinates q_i and momenta p_i in a single array w_μ).

Hint: $y_1 = Q$, $y_2 = \omega P$, where Q and P are the new generalized coordinates and momenta, respectively.

¹In 3 dimensions, the (antisymmetric) Levi-Civita tensor is defined as $\epsilon_{123} = \epsilon_{231} = \epsilon_{312} = 1$, $\epsilon_{132} = \epsilon_{213} = \epsilon_{321} = -1$, all other $\epsilon_{ijk} = 0$. In n dimensions $\epsilon_{123\dots n}$ and its even permutations (i.e., even number of swapping of adjacent indices) are 1, odd permutations -1 , all others 0.