


## Let's move to Bound States

When we discuss bound states of two objects in central-force potential, kinetic energy and potential energy are  $\sim$ the same. How does this compare to the rest energy of the objects?

Hydrogen ionization energy: 13.6 eV vs 0.5 MeV rest mass


Masses of b and c quarks are  $\sim$ relatively large, so we can also consider them non-relativistically (which makes them much easier). NOT true for uds quarks

# Briefest of overviews of hydrogen atom



Fine structure from relativistic corrections and spin-orbit coupling

Energy levels  $\sim -1/n^2$



$|s(e)| = 1/2$   
so  $j = l \pm 1/2$


Fig. 5.2 Fine structure in hydrogen. The  $n$ th Bohr level (fine line) splits into  $n$  sub-levels (dashed lines), characterized by  $j = \frac{1}{2}, \frac{3}{2}, \dots, (n - \frac{1}{2})$ . Except for the last of these, two different values of  $l$  contribute to

each level:  $l = j - \frac{1}{2}$  and  $l = j + \frac{1}{2}$ . Spectroscopists' nomenclature – S for  $l = 0$ , P for  $l = 1$ , D for  $l = 2$ , F for  $l = 3$  – is indicated. All levels are shifted downward, as shown (the diagram is not to scale, however).

Griffiths

# Lamb shift

Lamb shift: Led to development of quantum electro-dynamics! QED corrections to the electron-proton interaction break degeneracy of two levels with same  $n, j$  but different  $L$  (so  $2S_{1/2}$  and  $2P_{1/2}$  are not fully degenerate)




Vacuum polarization

Electron mass renormalization

Anomalous magnetic moment

Fig. 5.3 Some loop diagrams contributing to the Lamb shift.



Willis  
Lamb

## Finally, hyperfine splitting


Spin-orbit coupling is principally due to spin of electron interacting with 'B field' from nucleus (fine structure). Much smaller is spin of nucleus interacting with 'B field' from electron. Goes as  $(m_e/m_p)^4$  hence hyperfine (and not fine) splitting.

For  $n=1$ , the difference in energy states of proton (e and p spins aligned vs anti-aligned, which is lower) is  $5.9 \mu\text{eV} = 1420 \text{ MHz} = 21 \text{ cm}$ . Famous 21 cm line (penetrates dust!)

## On the 21 cm line

Lifetime of 21 cm is millions of years! Thankfully, enough hydrogen can provide this transition. Long lifetime = narrow width, so this is excellent for spectroscopy (Doppler shifts)

Used extensively in radio-astronomy, studying the early Universe, galaxy formation, measuring distances to objects, cosmology



Pioneer Plaque: 21 cm line defines distance and unit of time

# On to quarkonium

Differences between quarkonium and hydrogen/positronium: Don't really know the potential (strong force!) Also, interaction between quarks is large. Doesn't work for two light quarks, either

Instead of considering different states as energy levels of an atom, consider different bound states as different particles, each with a different mass. Start with mesons (much easier than baryons)



# What sorts of potentials can we use?



At short distances, we know that QCD is not a strong force.

Reasonable to start with  $1/r$  potential.

At large distances, we know that force grows exponentially. Try  $V \sim kr$  (others could also work). Of course,  $k$  can be a function of  $r$  too!

## Charmonium

Griffiths





Labeled (confusingly, to me!) as  $n^{(2s+1)}L_J$  ( $L=S,P,D,\dots$  for  $0,1,2,\dots$  and  $s=0$  or  $1$  for anti-aligned or aligned spins), with  $J=L+s$



## Charmonium

Griffiths



J/Psi state discovered in 1974

# Why was J/Psi discovered first?

$\eta$

$$J^{PC} = 0^+(0^-+)$$

From PDG

$\Gamma_i$	charged modes	(28.06 ± 0.34) %	S=1.2
$\Gamma_8$	charged modes	(28.06 ± 0.34) %	S=1.2
$\Gamma_9$	$\pi^+\pi^-\pi^0$	(22.73 ± 0.28) %	S=1.2
$\Gamma_{10}$	$\pi^+\pi^-\gamma$	( 4.60 ± 0.16 ) %	S=2.1
$\Gamma_{11}$	$e^+e^-\gamma$	( 6.8 ± 0.8 ) × 10 <sup>-3</sup>	S=1.7
$\Gamma_{12}$	$\mu^+\mu^-\gamma$	( 3.1 ± 0.4 ) × 10 <sup>-4</sup>	
$\Gamma_{13}$	$e^+e^-$	< 7.7 × 10 <sup>-5</sup>	CL=90%
$\Gamma_{14}$	$\mu^+\mu^-$	( 5.8 ± 0.8 ) × 10 <sup>-6</sup>	
$\Gamma_{15}$	$e^+e^-e^+e^-$	< 6.9 × 10 <sup>-5</sup>	CL=90%
$\Gamma_{16}$	$\pi^+\pi^-e^+e^-$	( 4.2 ± 1.2 ) × 10 <sup>-4</sup>	
$\Gamma_{17}$	$\pi^+\pi^-2\gamma$	< 2.0 × 10 <sup>-3</sup>	
$\Gamma_{18}$	$\pi^+\pi^-\pi^0\gamma$	< 5 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{19}$	$\pi^0\mu^+\mu^-\gamma$	< 3 × 10 <sup>-6</sup>	CL=90%

Due to C-parity!

$J/\psi(1S)$


$$J^{PC} = 0^-(1^{--})$$

## J/ψ(1S) DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ hadrons	(87.7 ± 0.5) %	
$\Gamma_2$ virtual $\gamma \rightarrow$ hadrons	(13.50 ± 0.30) %	
$\Gamma_3$ $ggg$	(64.1 ± 1.0) %	
$\Gamma_4$ $\gamma gg$	( 8.8 ± 0.5 ) %	
$\Gamma_5$ $e^+e^-$	( 5.94 ± 0.06 ) %	
$\Gamma_6$ $\mu^+\mu^-$	( 5.93 ± 0.06 ) %	

# Bottomonium

Griffiths



Even richer set of states!

Upsilon discovered in 1976

# How to account for spin in hadron masses?

Recall magnetic moment formula:  $\mu = -\frac{e}{m}\mathbf{S}$

Spin-spin interactions in hadrons  
have two components:

$$\mu_1 \cdot \mathbf{S}_2 = -\frac{e}{m_1}\mathbf{S}_1 \cdot \mathbf{S}_2$$

$$\mu_2 \cdot \mathbf{S}_1 = -\frac{e}{m_2}\mathbf{S}_2 \cdot \mathbf{S}_1 = -\frac{e}{m_2}\mathbf{S}_1 \cdot \mathbf{S}_2$$

Sum is then

$$-e\frac{m_1 + m_2}{m_1 m_2} (\mathbf{S}_1 \cdot \mathbf{S}_2) = A(m_1 + m_2)\frac{1}{m_1 m_2} (\mathbf{S}_1 \cdot \mathbf{S}_2)$$

# What is the spin term?

$$(\mathbf{S}_1 \cdot \mathbf{S}_2), \mathbf{S} = \mathbf{S}_1 + \mathbf{S}_2$$

$$\mathbf{S}^2 = \mathbf{S}_1^2 + \mathbf{S}_2^2 + 2(\mathbf{S}_1 \cdot \mathbf{S}_2)$$

$$(\mathbf{S}_1 \cdot \mathbf{S}_2) = \frac{1}{2} (\mathbf{S}^2 - \mathbf{S}_1^2 - \mathbf{S}_2^2)$$

$S_1$  and  $S_2 = \pm 1/2$

$$\mathbf{S}_1^2 = \mathbf{S}_2^2 = (1/2)(1/2+1) = 3/4$$

$$\mathbf{S}^2 = (1)(1+1) = 2 \text{ (spins aligned) or}$$

$$\mathbf{S}^2 = (0)(0+1) = 0 \text{ (spins anti-aligned)}$$

$$\text{So: } \mathbf{S}_1 \cdot \mathbf{S}_2 = 1/4 \text{ (spins aligned)}$$

$$\text{or: } \mathbf{S}_1 \cdot \mathbf{S}_2 = -3/4 \text{ (spins anti-aligned)}$$

# How to account for spin in hadron masses?

$$M(m_1, \dots, m_2) = m_1 + m_2 + A(m_1 + m_2) \frac{1}{m_1 m_2} (\mathbf{S}_1 \cdot \mathbf{S}_2)$$

Mass  $M$  of meson composed of quarks with mass  $m_1$  and  $m_2$  then generically looks like this

$$\mathbf{S}_1 \cdot \mathbf{S}_2 = 1/4 \text{ (spins aligned)}$$

$$\text{or: } \mathbf{S}_1 \cdot \mathbf{S}_2 = -3/4 \text{ (spins anti-aligned)}$$

Can try something simpler, and assume  $A$  is a constant (it surely is not, but maybe that is a reasonable approximation)

## Fits for masses (from Griffiths)

Table 5.3 Pseudoscalar and vector meson masses. (MeV/c<sup>2</sup>)

Meson	Calculated	Observed
$\pi$	139	138
$K$	487	496
$\eta$	561	548
$\rho$	775	776
$\omega$	775	783
$K^*$	892	894
$\phi$	1031	1020

Very nice agreement! But need to be careful...

For example: 
$$\eta = \frac{u\bar{u} + d\bar{d} - 2s\bar{s}}{\sqrt{6}}$$

## On to baryons

A lot more complicated - have three quarks, and thus three spins to add together. Most importantly, mesons are always composed of a quark and an anti-quark, ie never contain two of the same particle. In baryons, however (example: proton = uud), this no longer has to be true.

Regardless, though, baryons have half-integer spin (three quarks with  $s = \pm 1/2$  can combine to give  $s = \pm 1/2$  or  $\pm 3/2$  only)



# How to add three spins

To add three spins together, we first start by adding two of them together. Back to those C-G tables from the PDG ...

## Combining two $1/2 \times 1/2$ particles

$$\left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle = \left| 1 \ 1 \right\rangle$$

$$\left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle = \sqrt{\frac{1}{2}} \left| 1 \ 0 \right\rangle + \sqrt{\frac{1}{2}} \left| 0 \ 0 \right\rangle$$

$$\left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle = \sqrt{\frac{1}{2}} \left| 1 \ 0 \right\rangle - \sqrt{\frac{1}{2}} \left| 0 \ 0 \right\rangle$$

$$\left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle = \left| 1 \ -1 \right\rangle$$

1/2 x 1/2		1		
	+1	1	0	
+1/2 +1/2	1	0	0	
+1/2 -1/2	1/2	1/2	1	
-1/2 +1/2	1/2	-1/2	-1	
	-1/2 -1/2	1		

ex:  
J=1

ex:  
 $m_1, m_2 =$   
 $-1/2, +1/2$

# Let's rearrange

$$\left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle = |1 \ 1 \rangle$$

$$\left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle = \sqrt{\frac{1}{2}} |1 \ 0 \rangle + \sqrt{\frac{1}{2}} |0 \ 0 \rangle$$

$$\left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle = \sqrt{\frac{1}{2}} |1 \ 0 \rangle - \sqrt{\frac{1}{2}} |0 \ 0 \rangle$$

$$\left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle = |1 \ -1 \rangle$$

These are the  
easy ones

$$|1 \ 1 \rangle = \left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle$$

$$|1 \ -1 \rangle = \left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle$$

Let's rearrange (can also use tables for this)

$$\left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle = |1 \ 1 \rangle$$

$$\left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle = \sqrt{\frac{1}{2}} |1 \ 0 \rangle + \sqrt{\frac{1}{2}} |0 \ 0 \rangle$$

Add together here

$$\left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle = \sqrt{\frac{1}{2}} |1 \ 0 \rangle - \sqrt{\frac{1}{2}} |0 \ 0 \rangle$$

$$\left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle = |1 \ -1 \rangle$$

$$\sqrt{2} |1 \ 0 \rangle = \left( \left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle \right) + \left( \left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle \right)$$

$$|1 \ 0 \rangle = \frac{1}{\sqrt{2}} \left( \left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle \right) + \frac{1}{\sqrt{2}} \left( \left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle \right)$$

Let's rearrange (can also use tables for this)

$$\left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle = |1 \ 1 \rangle$$

$$\left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle = \sqrt{\frac{1}{2}} |1 \ 0 \rangle + \sqrt{\frac{1}{2}} |0 \ 0 \rangle$$

Subtract these

$$\left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle = \sqrt{\frac{1}{2}} |1 \ 0 \rangle - \sqrt{\frac{1}{2}} |0 \ 0 \rangle$$

$$\left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle = |1 \ -1 \rangle$$

$$\sqrt{2} |0 \ 0 \rangle = \left( \left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle \right) - \left( \left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle \right)$$

$$|0 \ 0 \rangle = \frac{1}{\sqrt{2}} \left( \left| \frac{1}{2} \frac{1}{2} \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle \right) - \frac{1}{\sqrt{2}} \left( \left| \frac{1}{2} \frac{-1}{2} \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle \right)$$

# Putting it together

$$|0\ 0\rangle = \frac{1}{\sqrt{2}} \left( \left| \frac{1}{2}\ \frac{1}{2} \right\rangle + \left| \frac{1}{2}\ \frac{-1}{2} \right\rangle \right) - \frac{1}{\sqrt{2}} \left( \left| \frac{1}{2}\ \frac{-1}{2} \right\rangle + \left| \frac{1}{2}\ \frac{1}{2} \right\rangle \right)$$

$$|1\ 0\rangle = \frac{1}{\sqrt{2}} \left( \left| \frac{1}{2}\ \frac{1}{2} \right\rangle + \left| \frac{1}{2}\ \frac{-1}{2} \right\rangle \right) + \frac{1}{\sqrt{2}} \left( \left| \frac{1}{2}\ \frac{-1}{2} \right\rangle + \left| \frac{1}{2}\ \frac{1}{2} \right\rangle \right)$$

$$|1\ 1\rangle = \left| \frac{1}{2}\ \frac{1}{2} \right\rangle + \left| \frac{1}{2}\ \frac{1}{2} \right\rangle$$

$$|1\ -1\rangle = \left| \frac{1}{2}\ \frac{-1}{2} \right\rangle + \left| \frac{1}{2}\ \frac{-1}{2} \right\rangle$$

When we add the third quark we will have to add spin 1/2 with either spin 0 or spin 1

# Now we add the third one

## Combining spin 1 x 1/2 particles

We want the “inverse” of what we have been reading off. Can also use the tables for that!

ex:  
 $J=3/2$   
 $m=1/2$

ex:  $m_1, m_2 = +1, -1/2$

$1 \times 1/2$	$3/2$	$3/2$	$1/2$		
$+1$	$+3/2$	$1$	$+1/2$	$+1/2$	
$+1$	$-1/2$	$1/3$	$2/3$	$3/2$	$1/2$
$0$	$+1/2$	$2/3$	$-1/3$	$-1/2$	$-1/2$
		$0$	$-1/2$	$2/3$	$1/3$
		$-1$	$+1/2$	$1/3$	$-2/3$
				$-1$	$-1/2$
					$1$

# Now we add the third one

## Combining spin 1 x 1/2 particles

$$|\frac{3}{2} \frac{3}{2}\rangle = |1 \ 1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle$$

$$|\frac{3}{2} \ \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} \left( |1 \ 1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) + \sqrt{\frac{2}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$


$$|\frac{1}{2} \ \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} \left( |1 \ 1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) - \sqrt{\frac{1}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{3}{2} \ \frac{-1}{2}\rangle = \sqrt{\frac{2}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) + \sqrt{\frac{1}{3}} \left( |1 \ -1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{1}{2} \ \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) - \sqrt{\frac{2}{3}} \left( |1 \ -1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{3}{2} \ \frac{-3}{2}\rangle = |1 \ -1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle$$

ex:  
J=3/2  
m=1/2



Combining spin 0 x 1/2 particles is trivial

$$\begin{aligned} \left| \frac{1}{2} \frac{1}{2} \right\rangle &= \left| 0 \ 0 \right\rangle + \left| \frac{1}{2} \frac{1}{2} \right\rangle \\ \left| \frac{1}{2} \frac{-1}{2} \right\rangle &= \left| 0 \ 0 \right\rangle + \left| \frac{1}{2} \frac{-1}{2} \right\rangle \end{aligned}$$



## So in total

$$|\frac{3}{2} \frac{3}{2}\rangle = |1 \ 1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle$$

$$|\frac{3}{2} \ \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} \left( |1 \ 1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) + \sqrt{\frac{2}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{1}{2} \ \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} \left( |1 \ 1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) - \sqrt{\frac{1}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{3}{2} \ \frac{-1}{2}\rangle = \sqrt{\frac{2}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) + \sqrt{\frac{1}{3}} \left( |1 \ -1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{1}{2} \ \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) - \sqrt{\frac{2}{3}} \left( |1 \ -1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{3}{2} \ \frac{-3}{2}\rangle = |1 \ -1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle$$

$$\begin{aligned} |0 \ 0\rangle &= \frac{1}{\sqrt{2}} \left( |\frac{1}{2} \ \frac{1}{2}\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) - \frac{1}{\sqrt{2}} \left( |\frac{1}{2} \ \frac{-1}{2}\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right) \\ |1 \ 0\rangle &= \frac{1}{\sqrt{2}} \left( |\frac{1}{2} \ \frac{1}{2}\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) + \frac{1}{\sqrt{2}} \left( |\frac{1}{2} \ \frac{-1}{2}\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right) \end{aligned}$$

$$|1 \ 1\rangle = |\frac{1}{2} \ \frac{1}{2}\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle$$

$$|1 \ -1\rangle = |\frac{1}{2} \ \frac{-1}{2}\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle$$

From first  
combination

$$|0 \ 0\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle = |\frac{1}{2} \ \frac{1}{2}\rangle$$

$$|0 \ 0\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle = |\frac{1}{2} \ \frac{-1}{2}\rangle$$

# Let's introduce some nicer notation

$$|\frac{3}{2} \frac{3}{2}\rangle = |1 \ 1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle$$

$$|\frac{3}{2} \ \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} \left( |1 \ 1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) + \sqrt{\frac{2}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{1}{2} \ \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} \left( |1 \ 1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) - \sqrt{\frac{1}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{3}{2} \ \frac{-1}{2}\rangle = \sqrt{\frac{2}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) + \sqrt{\frac{1}{3}} \left( |1 \ -1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{1}{2} \ \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( |1 \ 0\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) - \sqrt{\frac{2}{3}} \left( |1 \ -1\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|\frac{3}{2} \ \frac{-3}{2}\rangle = |1 \ -1\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle$$

$$|0 \ 0\rangle = \frac{1}{\sqrt{2}} \left( |\frac{1}{2} \ \frac{1}{2}\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) - \frac{1}{\sqrt{2}} \left( |\frac{1}{2} \ \frac{-1}{2}\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|1 \ 0\rangle = \frac{1}{\sqrt{2}} \left( |\frac{1}{2} \ \frac{1}{2}\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle \right) + \frac{1}{\sqrt{2}} \left( |\frac{1}{2} \ \frac{-1}{2}\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle \right)$$

$$|1 \ 1\rangle = |\frac{1}{2} \ \frac{1}{2}\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle$$

$$|1 \ -1\rangle = |\frac{1}{2} \ \frac{-1}{2}\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle$$

From first  
combination

Don't forget: order matters!

$$|0 \ 0\rangle + |\frac{1}{2} \ \frac{1}{2}\rangle = |\frac{1}{2} \ \frac{1}{2}\rangle$$

$$|0 \ 0\rangle + |\frac{1}{2} \ \frac{-1}{2}\rangle = |\frac{1}{2} \ \frac{-1}{2}\rangle$$

$$|\frac{1}{2} \ \frac{1}{2}\rangle = (\uparrow)$$

$$|\frac{1}{2} \ \frac{-1}{2}\rangle = (\downarrow)$$

## Using the notation

$$|\frac{3}{2} \frac{3}{2}\rangle = |1 1\rangle + (\uparrow)$$

$$|\frac{3}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} (|1 1\rangle + (\downarrow)) + \sqrt{\frac{2}{3}} (|1 0\rangle + (\uparrow))$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} (|1 1\rangle + (\downarrow)) - \sqrt{\frac{1}{3}} (|1 0\rangle + (\uparrow))$$

$$|\frac{3}{2} \frac{-1}{2}\rangle = \sqrt{\frac{2}{3}} (|1 0\rangle + (\downarrow)) + \sqrt{\frac{1}{3}} (|1 -1\rangle + (\uparrow))$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} (|1 0\rangle + (\downarrow)) - \sqrt{\frac{2}{3}} (|1 -1\rangle + (\uparrow))$$

$$|\frac{3}{2} \frac{-3}{2}\rangle = |1 -1\rangle + (\downarrow)$$

$$|\frac{1}{2} \frac{1}{2}\rangle = |0 0\rangle + \uparrow$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = |0 0\rangle + \downarrow$$

$$|0 0\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow) - \frac{1}{\sqrt{2}} (\downarrow\uparrow)$$

$$|1 0\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow) + \frac{1}{\sqrt{2}} (\downarrow\uparrow)$$

$$|1 1\rangle = \uparrow\uparrow$$

$$|1 -1\rangle = \downarrow\downarrow$$

# Putting it together

$$|\frac{3}{2} \frac{3}{2}\rangle = \uparrow\uparrow\uparrow$$

$$|\frac{3}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} (\uparrow\uparrow\downarrow) + \sqrt{\frac{2}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} (\uparrow\uparrow\downarrow) - \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{3}{2} \frac{-1}{2}\rangle = \sqrt{\frac{2}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) (\downarrow) \right) + \sqrt{\frac{1}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \downarrow \right) - \sqrt{\frac{2}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{3}{2} \frac{-3}{2}\rangle = \downarrow\downarrow\downarrow$$

Phew

$$|\frac{1}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) \uparrow$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) \downarrow$$

# How to interpret

$$|\frac{3}{2} \frac{3}{2}\rangle = \uparrow\uparrow\uparrow$$

$$|\frac{3}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} (\uparrow\uparrow\downarrow) + \sqrt{\frac{2}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} (\uparrow\uparrow\downarrow) - \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{3}{2} \frac{-1}{2}\rangle = \sqrt{\frac{2}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) (\downarrow) \right) + \sqrt{\frac{1}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \downarrow \right) - \sqrt{\frac{2}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{3}{2} \frac{-3}{2}\rangle = \downarrow\downarrow\downarrow$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) \uparrow$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) \downarrow$$

$$|\frac{3}{2} \frac{3}{2}\rangle = \uparrow\uparrow\uparrow$$

$$|\frac{3}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} (\uparrow\uparrow\downarrow + \uparrow\downarrow\uparrow + \downarrow\uparrow\uparrow)$$

$$|\frac{3}{2} \frac{-1}{2}\rangle = \frac{1}{\sqrt{3}} (\uparrow\downarrow\downarrow + \downarrow\uparrow\downarrow + \downarrow\downarrow\uparrow)$$

$$|\frac{3}{2} \frac{-3}{2}\rangle = \downarrow\downarrow\downarrow$$

Spin 3/2 states  
are easy to  
interpret:  
symmetric if we  
interchange any  
two quarks

# How to interpret

$$|\frac{3}{2} \frac{3}{2}\rangle = \uparrow\uparrow\uparrow$$

$$|\frac{3}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} (\uparrow\uparrow\downarrow) + \sqrt{\frac{2}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} (\uparrow\uparrow\downarrow) - \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{3}{2} \frac{-1}{2}\rangle = \sqrt{\frac{2}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) (\downarrow) \right) + \sqrt{\frac{1}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \downarrow \right) - \sqrt{\frac{2}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{3}{2} \frac{-3}{2}\rangle = \downarrow\downarrow\downarrow$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} (\uparrow\uparrow\downarrow) - \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \downarrow \right) - \sqrt{\frac{2}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) \uparrow$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) \downarrow$$

Two of spin 1/2 states are asymmetric under interchange of first and second quarks

# How to interpret

$$|\frac{3}{2} \frac{3}{2}\rangle = \uparrow\uparrow\uparrow$$

$$|\frac{3}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{3}} (\uparrow\uparrow\downarrow) + \sqrt{\frac{2}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} (\uparrow\uparrow\downarrow) - \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{3}{2} \frac{-1}{2}\rangle = \sqrt{\frac{2}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) (\downarrow) \right) + \sqrt{\frac{1}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \downarrow \right) - \sqrt{\frac{2}{3}} (\downarrow\downarrow\uparrow)$$

$$|\frac{3}{2} \frac{-3}{2}\rangle = \downarrow\downarrow\downarrow$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) \uparrow$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \frac{1}{\sqrt{2}} (\uparrow\downarrow - \downarrow\uparrow) \downarrow$$

$$|\frac{1}{2} \frac{1}{2}\rangle = \sqrt{\frac{2}{3}} (\uparrow\uparrow\downarrow) - \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \uparrow \right)$$

$$|\frac{1}{2} \frac{-1}{2}\rangle = \sqrt{\frac{1}{3}} \left( \frac{1}{\sqrt{2}} (\uparrow\downarrow + \downarrow\uparrow) \downarrow \right) - \sqrt{\frac{2}{3}} (\downarrow\downarrow\uparrow)$$

These last two spin 1/2 states are symmetric under interchange of first and second quarks

We need our 3 quarks to satisfy Fermi-Dirac statistics (must be anti-symmetric under exchange of any two quarks)

For ground state ( $l=0$ ), space wave function is symmetric. Left off with wave functions for **spin**, **color** and **flavor**. We will see that **color** wave function is necessarily anti-symmetric. That means that **flavor x spin** combination must be symmetric



You don't have Chapter 5 HW, but ...

## Something new to think about

I'd like for you to begin to think about your final project. You should put in some effort to prepare it, so time to start now. You need to pick a single **analysis** or result to present (something public in the past ~year)

<http://arxiv.org/archive/hep-ex>

Minos  
ATLAS  
CMS  
LHCb  
ALICE  
SNO  
Belle-2  
AMS  
LIGO  
VIRGO  
LUX  
Dark Energy Survey  
Pierre Auger  
X17  
Fermi Gamma Ray  
Telescope

Pick an analysis/paper and email it to me for approval no later than 1 week from today. I'll ask for some progress reports in the future, but for now I just want you to pick something interesting (and not what you work on for research, and not what someone else will be talking about). Talk to me if you need help picking a topic!

## Some more news on your presentation

Want a 25 minute presentation on the topic! You should be including theory background if possible, as well as information on the detector, the analysis, the background estimation, and the significance of the result. We will all discuss the presentation for 10 minutes after you're done (aka ask you some questions)

**I also reserve the right to reject papers that are too broad, too narrow, too old, or too out of focus for this course**

## Aside on your final presentation

Note 0: It's not first-come/first-served, but instead we will flip coins or play rock-paper-scissors for who gets which topic.

Note 1: If you dropped by my office to discuss the paper, that does not count as fulfilling your homework assignment. Please sent it to me by email

Note 2: This really counts as a homework assignment. So don't miss the deadline! If you do, you get points off - and I get to pick a topic for you :)

## Aside on your final presentation

Note 3: Only one topic per person, so you might want to have 1-2 backups in mind (class is small so maybe that's OK). I'll let you know class after it's due where we have duplicates (and alternates due then class after that)

Note 4: You are not covering an “experiment”, but rather a single analysis/limit/measurement. So I want you to have the paper you will be reading in mind. If there is no physics result, then this does not count