

Pitch correction


D. Rubin

USPAS January 2019

Suppose the muon is oscillating in a vertical plane at the magic radius

$$\psi = \psi_0 \sin \omega_p t$$

$$\omega_a = \frac{ae}{mc\gamma} B^* \quad \text{In the rest frame of the muon}$$



$$\mathbf{B}' = \gamma \mathbf{B} - \frac{\gamma^2}{\gamma + 1} \beta (\beta \cdot \mathbf{B})$$

$$B_z^{eff} = B_z \left(1 - \frac{\gamma - 1}{\gamma} \psi^2\right) \quad B_x = -B_z \left(\frac{\gamma - 1}{\gamma} \psi\right)$$

$$\omega_z = \omega_0 \left(1 - \frac{\gamma - 1}{\gamma} \psi^2\right) \quad \omega_x = -\omega_0 \left(\frac{\gamma - 1}{\gamma} \psi\right)$$

Where $\omega_0 = a_\mu (e/mc) B_z$

The vertical restoring force (E or B) provides a third axes

$$\omega_y = -f \omega_p \psi_0 \cos \omega_p t \quad \text{where} \quad f = (1 + \beta^2 \gamma a - \gamma^{-1})$$

$$x = \cos \theta \cos \phi$$

$$y = \cos \theta \sin \phi$$

$$z = \sin \theta$$

$$\dot{\theta} = \omega_x \sin \phi - \omega_y \cos \phi = -\omega_0 \psi_0 \frac{\gamma - 1}{\gamma} \sin \phi \sin \omega_p t + \omega_p \psi_0 f \cos \phi \cos \omega_p t$$

$$\begin{aligned} \dot{\phi} &= \omega_z - \omega_x \tan \theta \cos \phi - \omega_y \tan \theta \sin \phi \\ &= \omega_0 \left(1 - \frac{\gamma - 1}{\gamma} \psi_0^2 \sin^2 \omega_p t\right) + \omega_0 \psi_0 \frac{\gamma - 1}{\gamma} \tan \theta \cos \phi \sin \omega_p t + \omega_p \psi_0 f \tan \theta \sin \phi \cos \omega_p t \end{aligned}$$

Substitute the lowest order time dependence for $\phi = (\omega t + \xi)$

And integrate

$$\theta = \frac{A_1}{\omega_0 + \omega_p} \sin((\omega_0 + \omega_p)t + \xi) - \frac{A_2}{\omega_0 - \omega_p} \sin((\omega_0 - \omega_p)t + \xi)$$

$$\dot{\phi} = \omega_0 \left(1 - \frac{\gamma - 1}{2\gamma} \psi_0^2 \right) + \frac{A_1^2}{2(\omega_0 + \omega_p)} + \frac{A_2^2}{2(\omega_0 - \omega_p)}$$

$$A_1 = \frac{1}{2} \psi_0 (\omega_0 \gamma^{-1} (\gamma - 1) + f \omega_p)$$

$$A_2 = \frac{1}{2} \psi_0 (\omega_0 \gamma^{-1} (\gamma - 1) - f \omega_p)$$

$$\dot{\phi} = \omega_0 \left(1 - \frac{\gamma - 1}{2\gamma} \psi_0^2 \right) + \frac{A_1^2}{2(\omega_0 + \omega_p)} + \frac{A_2^2}{2(\omega_0 - \omega_p)}$$

This is what we measure

At the magic momentum

$$\dot{\phi} = \omega_0(1 - C)$$

$$C = \frac{1}{4} \psi_0^2 (1 - (\omega_0^2 + 2\pi\gamma^2) / \gamma^2 (\omega_0^2 - \omega_p^2))$$

Resonance when $\omega_p \sim \omega_0$

Otherwise $C \sim \frac{1}{4} \psi_0^2$

Assume that polarization is in the plane

where ω_v is the vertical betatron frequency and ψ is the pitch angle. Define θ so that $\hat{\beta} \cdot \mathbf{s} = s \cos \theta$ and $\hat{\beta} \times \mathbf{s} = s \sin \theta \hat{\mathbf{s}}_{\perp}$. Then $\hat{\mathbf{s}}_{\perp} \cdot \mathbf{B} = B \cos \psi$. (B in the vertical direction). Now

$$\begin{aligned} \frac{d(\hat{\beta} \cdot \mathbf{s})}{dt} &= -\frac{e}{mc} \mathbf{s}_{\perp} \cdot \left[\left(\frac{g}{2} - 1 \right) (\hat{\beta} \times \mathbf{B}) + \left(\frac{g}{2} - 1 - \frac{1}{\gamma^2 - 1} \right) \mathbf{E} \right] \\ \frac{sd(\cos \theta)}{dt} &= -\frac{e}{mc} \mathbf{B} \cdot \left(\frac{g}{2} - 1 \right) (\mathbf{s} \times \hat{\beta}) \\ -\sin \theta \frac{d\theta}{dt} &= -\frac{e}{mc} B \left(\frac{g}{2} - 1 \right) \cos \psi (s \sin \theta) \\ \rightarrow \omega_a &= -\frac{e}{mc} B \left(\frac{g}{2} - 1 \right) \cos \psi \end{aligned}$$

Then $\beta \times \mathbf{B} = \hat{\mathbf{i}} \psi B \cos \omega_v t$

$\hat{\beta} = (\cos \alpha \hat{\mathbf{k}} + \sin \alpha \hat{\mathbf{j}})$. Note that $\mathbf{s}_{\perp} \cdot (\hat{\beta} \times \mathbf{B}) = \mathbf{s} \cdot (\hat{\beta} \times \mathbf{B})$. Then 5 becomes

$$\frac{d(\hat{\beta} \cdot \mathbf{s})}{dt} = \frac{d}{dt} (s(\sin \alpha \sin \theta \sin \phi + \cos \alpha \cos \theta)) \sim \frac{d}{dt} (s(\cos \alpha \cos \theta)) \quad (10)$$

The pitch angle α oscillates with the vertical betatron frequency which is much greater than the precession frequency with average value of zero. Then

$$\begin{aligned} \frac{d(\hat{\beta} \cdot \mathbf{s})}{dt} &= -s(\cos \alpha \sin \theta) \frac{d\theta}{dt} = sB \sin \theta \cos \phi \cos \alpha \left(\frac{g}{2} - 1 \right) \\ \rightarrow \omega &= -B \cos \phi \left(\frac{g}{2} - 1 \right) \end{aligned}$$

Pedestrian estimate of pitch correction

$$\frac{d}{dt}(\hat{\beta} \cdot \mathbf{s}) = -\frac{e}{mc} \mathbf{s}_{\perp} \cdot \left[\left(\frac{g}{2} - 1 \right) \hat{\beta} \times \mathbf{B} + \left(\frac{g\beta}{2} - \frac{1}{\beta} \right) \mathbf{E} \right]$$

$$\hat{\beta} \cdot \mathbf{s} = \cos \phi \qquad \hat{\beta} \times \mathbf{B} = B \cos \psi \hat{x}$$

$$\frac{d}{dt} \cos \phi = -\frac{e}{mc} \sin \phi (a_{\mu} B) \cos \psi$$

$$\dot{\phi} = -\frac{e}{mc} a_{\mu} B \left(1 - \frac{1}{2} \psi^2 \right)$$



$$= \omega_0 \left(1 - \frac{1}{2} \psi^2 \right)$$

$$\rightarrow C = -\frac{1}{2} \langle \psi^2 \rangle = -\frac{1}{4} \psi_0^2$$

Pitch correction

$$\vec{C}_p(T) = \frac{1}{B} \frac{1}{T} \int_0^T (\vec{\beta} \cdot \vec{B}) \vec{\beta} dt$$

1. Generated and track a distribution and compute
 - Momentum and frequency distribution
 - E-field correction for each particle
 - Vertical phase space distribution and pitch correction



on momentum, pitch

